

DORIN, V.A.; KUZNETSOV, B.I.; NASLEDOV, D.N.

Investigating the growth of a layer of an n-type semiconductor at
a cadmium-selenium contact. *Fiz.tver.tela* 1 no.5:734-739 My '59.
(MIRA 12:4)

1. Leningradskiy fiziko-tekhnicheskii institut AN SSSR.
(Cadmium) (Selenium) (Semiconductors)

5(2)

AUTHORS:

Dorin, V. A., Tartakovskaya, F. M.

05889

SOV/78-4-11-42/50

TITLE:

The Reduction of Titanium Dioxide in the Presence of Titanium

PERIODICAL:

Zhurnal neorganicheskoy khimii, 1959, Vol 4, Nr 11,
pp 2635-2637 (USSR)

ABSTRACT:

The reduction of TiO_2 has so far always been carried out by means of direct contact of the reagent with TiO_2 . In the present paper, the authors report on the reduction of TiO_2 by means of Ti without contact between the two substances. Ti, in a quartz container, was submerged into the quartz test glass filled with TiO_2 so that the reduction could only take place by way of the gaseous phase. The behavior of the TiO_2 -modifications rutile and anatase was investigated at temperatures up to 1100° (Tables 1,2). The color changes observed at rising temperature are caused by Ti^{3+} -ions. After heating for five hours, the rutile had the composition $TiO_{1.936}$. Traces of Ti_3O_5 appeared at 1050° .

Card 1/2

The Reduction of Titanium Dioxide in the
Presence of Titanium

05889

SOV/78-4-11-42/50

The anatase was transformed into rutile. The reduction of TiO_2 in the presence of Ti takes place within a wide temperature range. By corresponding variation in temperature and reaction time, dioxides with any deviation from the stoichiometric ratio can be obtained. Here, the TiO_2 becomes a semiconductor. The authors thank D. N. Nasledov for the attention paid to the paper. There are 2 tables and 5 references.

ASSOCIATION: Leningradskiy fiziko-tekhnicheskii institut Akademii nauk SSSR
(Leningrad Physical-technical Institute of the Academy of
Sciences, USSR)

SUBMITTED: April 22, 1959

Card 2/2

68022

S/126/60/009/02/007/033

E111/E335

5.2200

AUTHORS:

Dorin, V.A. and Filaretova, G.M.

TITLE:

Investigation of the Growth of a Lead Sulphide Film on
Lead in Contact with Liquid Sulphur

PERIODICAL:

Fizika metallov i metallovedeniye, 1960, Vol 9, Nr 2,
pp 195 - 201 (USSR)

ABSTRACT:

The process of growth was studied in the range 175 to 300 °C by placing lead samples in liquid sulphur. X-ray analysis showed that the film formed consisted of one compound only with the composition of PbS.¹ Figure 1 shows a micrograph of the sulphide film obtained at 200 °C. the thickness was measured by a comparator IZA-2. Figure 2 shows that there is a linear relationship between the thickness and time. The rate of growth has an exponential relationship with temperature (Figure 3). The effect of impurities, in both the lead and sulphur was studied.. The films obtained on lead containing 1 at. % Sn, Cd or Zn were very thin. An addition of 1 at. % of Cu had no effect. Experiments in vacuo showed that the presence of air considerably ✓

Card1/2

68622

S/126/60/009/02/007/033

E111/E335

Investigation of the Growth of a Lead Sulphide Film on Lead in
Contact with Liquid Sulphur

increased the thickness of the film. It was shown that oxygen and not nitrogen was the cause of this increase. The presence of selenium or tellurium in the sulphur also gave a marked increase in growth. It was demonstrated that the growth of lead sulphide took place at the lead-lead sulphide interface. The rate of growth of the film is determined not by the diffusion through the film but by the rate of formation of lead sulphide.

Acknowledgments are expressed to Professor D.N. Nasledov for his continued interest and for his comments on the results. There are 5 figures, 2 tables and 12 references, 4 of which are English, 3 German and 5 Soviet.

ASSOCIATION: Leningradskiy fiziko-tekhnicheskii institut AN SSSR
(Leningrad Physico-technical Institute of the Ac.Sc., USSR)

SUBMITTED: July 13, 1959

Card 2/2

80531

18.7520

S/126/60/009/05/011/025

EQ21/E335

AUTHORS: Dorin, V.A. and Filaretova, G.M.

TITLE: The Growth of a Lead Selenide Film

PERIODICAL: Fizika metallov i metallovedeniye, 1960, Vol 9, Nr 5,
pp 718 - 721 (USSR)

ABSTRACT: Experiments were carried out on the diffusion of liquid selenium in contact with solid lead. The apparatus used (Figure 1) ensured that no diffusion could take place in the time taken to heat up to the experimental temperature. The apparatus was evacuated to 10^{-5} mm Hg and placed in a thermostat. The diffusion layer formed after several minutes consisted of two parts. Figure 2 shows the selenium 1, a porous PbSe layer 2, a compact PbSe layer 3 and lead 4. X-ray analysis showed that both the diffusion layers contained PbSe. Figure 3 shows the structure of the porous layer which consists of a network of lead selenide crystals, the pores of which are filled with amorphous selenium. Microhardness measurements confirmed this, giving values of 75 kg/mm^2 for PbSe and 57 kg/mm^2 for Se. The thickness of the compact layer

Card1/3

80531

S/126/60/009/05/011/025

E021/E335

The Growth of a Lead Selenide Film

was always the same under the same conditions but the thickness of the porous layer varied even when prepared under exactly the same conditions. With increase in time of diffusion, the compact layer increased in thickness and the porous layer decreased. A similar picture was obtained with increase in temperature. Oxygen had a pronounced effect on diffusion. The layer produced with the apparatus filled with air was several times thicker than that produced in vacuo. Only a thin compact layer (and no porous layer) is formed with solid selenium in contact with lead. The mechanisms of film formation with solid and liquid selenium are obviously different. Lead atoms diffuse into the liquid selenium to give the porous layer. This was confirmed by carrying out tests with lead covered with lead sulphide.

There are 4 figures and 6 references, 1 of which is English and 5 are Soviet.

4

Card2/3

80531

The Growth of a Lead Selenide Film

S/126/60/009/05/011/025
E021/E335

ASSOCIATION: Fiziko-tekhnicheskii institut AN SSSR, Leningrad
(Physics-engineering Institute of the Ac.Sc., USSR,
Leningrad)

SUBMITTED: July 13, 1959 - initially;
December 7, 1959 - after revision.

Card 3/3

✓

80227

8/076/60/034/04/18/042
B010/B009

5.2100

AUTHORS: Dorin, V. A., Nasledov, D. N., Tartakovskaya, F. M. (Leningrad)
TITLE: Preparation of a Titanium Dioxide Semiconductor on Titanium at Low Oxygen Pressures

PERIODICAL: Zhurnal fizicheskoy khimii, 1960, Vol. 34, No. 4, pp. 809 - 814

TEXT: The oxidation of titanium in a gaseous phase obtained by heating powdered titanium oxide was investigated. In this way a gaseous phase containing only small amounts of oxygen was obtained. Titanium foils (0.6 mm thick, 20 X 20 mm²) with at most 0.08% C, 0.08% N₂, 0.5% Fe + Ni, and traces of Cu were oxidized. The titanium oxide powder was annealed at 800° for three hours prior to use. In the first series of experiments anatase powder was used, in the second, rutile powder. Working temperatures ranged from 700° to 1100°, the weight increase in the titanium foil undergoing oxidation was determined by weighing. In the first series of experiments the color of the oxide film was observed to change with temperature, i.e., at 650-800° the oxide is light gray, but changes into dark gray and, at temperatures above 850°, into dark blue. An X-ray analysis showed that at

Card 1/2

Preparation of a Titanium Dioxide Semiconductor on Titanium at Low Oxygen Pressures

80227

S/076/60/034/04/18/042
B010/B009

temperatures up to 850-900° an oxide film with a rutile structure forms. At 1100° two oxide layers were found, namely a thin upper layer of Ti_3O_5 and a lower layer the X-ray picture of which was different, although its composition is likewise Ti_3O_5 . The dependence of the growth of the oxide layer upon time was found to be parabolic, while the temperature dependence is governed by an exponential law. The results of the second series of experiments (Table) show that the sample weight increases at 700-900° only. The oxidation of titanium takes place while the titanium dioxide powder is greatly reduced. The oxide film forming during the process has an electrical conductivity of the electronic type. This electrical conductivity depends on the temperature at which the oxide film is produced. G. P. Luchkin and G. G. Il'in are mentioned in the text. There are 5 figures, 1 table, and 19 references, 4 of which are Soviet.

SUBMITTED: June 27, 1958

✓

Card 2/2

BAKAYEV, A.V.; GELLER, I. Kh.; DORIN, V.A.; ZAKHAROV, M.P.; NASLEDOV, D.N.;
SOLOV'YEV, R.A.

Method for investigating potential distribution in selenium
rectifying cells. Zav.lab. 27 no.10:1240-1242 '61. (MIRA 14:10)

1. Leningradskiy politekhnicheskoy institut im. M. I. Kalinina.
(Selenium—Electric properties)

S/139/63/000/001/012/027
E202/E420

AUTHORS: Bakayev, A.V., Gellor, I.Kh., Dorin, V.A., Zakharov, P.M.,
Nasledov, D.N., Solov'yev, R.A.

TITLE: Distribution of potential in selenium rectifying
elements between electrodes

PERIODICAL: Izvestiya vysshikh uchobnykh zavedeniy. Fizika,
no.1, 1963, 78-84

TEXT: Results of measuring potential distribution in selenium
rectifying elements in the conducting direction are described.
To explain in detail the mechanism of potential distribution between
the electrodes, measurements were carried out at points separated
by a distance of 5μ . Since the thickness of selenium layer varies
from 50 to 100μ it was necessary to measure the potential at 10 to
20 points. In order to carry out the measurements the layer of
selenium and the p-n junction region were stripped and a transverse
section prepared. Both types of rectifiers, i.e. those with p-n
junction between the upper electrode and the layer of selenium,
and those in which the p-n junction lies between the layer of
selenium and the base, were investigated. The method was based on
Card 1/3

S/139/63/000/001/012/027
E202/E420

Distribution of potential ...

measuring the difference of potential between one of the electrodes and a probe, the latter being placed at various points on the surface of the transverse section of the element. A special instrument incorporating a microhardness gauge of the diamond pyramid type in which the latter was replaced by a steel wedge-shaped probe was used. During measurements the probe was pressed into the selenium in order to obtain reliable results. The width of the indentation made by the probe was 1.5 to 2 μ , hence the potential could be measured at points separated by a distance of 5 μ . Since the probe contact with selenium has a considerable resistance of the order of 10^8 to 10^9 ohms, a high resistance voltmeter was used in the measurements. This comprised a potentiometer with a center zero electrometer sensitive to a current of 10^{-11} A. The measurements had an absolute error of 0.001 V. Considerable care was taken in the preparation of the transverse sections. The results have shown that the main fraction of the potential applied to the element in the conducting direction falls over the p-n junction region, on the other hand the layer of selenium accounts for not more than 25% of the above fall. In addition to plotting

Card 2/3

Distribution of potential ...

S/139/63/000/001/012/027
E202/E420

the potential against the distance over the CdS-(orCdSe)-Se-Pi₂Se₃-Al portions of the sandwich, preliminary volt-ampere characteristics of both types of rectifier were measured on polished and unpolished samples. There are 6 figures.

ASSOCIATION: Leningradskiy politekhnicheskii institut imeni M.I.Kalinina (Leningrad Polytechnic Institute imeni M.I.Kalinin)

SUBMITTED: August 22, 1961

Card 3/3

L 14535-63

WP(q)/EWT(m)/BDS AFFTC/ASD JD

ACCESSION NR: AP3005310

8/0181/63/005/008/2065/2069

AUTHOR: Dorin, V. A.; Patrakova, A. Ya.

TITLE: Investigation of electrical characteristics of TiO_{2-x} -Ag contact with an intermediate insulating layer

SOURCE: Fizika tverdogo tela, v. 7, no. 8, 1963, 2065-2069

TOPIC TAGS: electrical-contact characteristic, electrical characteristic, contact characteristic, titanium oxide, titanium-oxide contact characteristic

ABSTRACT: The electrical characteristics of a TiO_{2-x} -Ag contact with an insulating layer have been investigated at room temperature and at 400C. The characteristics obtained were compared with those of a system without an insulating layer. It was found that unipolar conductivity exists up to 500C in specimens with insulating layers. Rectification takes place in the area of contact with the intermediate layer at 400C. At 300C the specimen had equal conductivity in both directions. In the region of low voltages the volt-ampere characteristics of the specimens indicated that the introduction of an insulating layer

Card 1/2

L 14535-63

ACCESSION NR: AP3005310

at room temperature only increases the resistance of the system. When a system with an insulating layer is heated to 400C the rectification factor is 30 at 100 mv. An analysis of separate sections of the volt-ampere characteristics in the reverse direction showed that the current increased proportionally with the increase of voltage up to approximately 1 v. Above 4 v, an exponential dependence takes place either at low or high temperature. The same dependence was observed in the contact area of Ag-TiO_{2-x} at lower voltages. Thus, the introduction of an insulating layer consisting of a material different from TiO_{2-x} does not change the qualitative picture of the characteristics. It serves only to elevate the temperature (by 200-250C) at which rectification takes place. Orig. art. has: 5 figures, 5 formulas, and 2 tables.

ASSOCIATION: Fiziko-tekhnicheskiy institut im. A. F. Ioffe AN SSSR, Leningrad
(Leningrad Physicotechnical Institute, AN SSSR)

SUBMITTED: 28Jan63

DATE ACQ: 06Sep63

ENCL: 00

SUB CODE: PH

NO REF SOV: 006

OTHER: 003

Card 2/2

DORIN, V.A.; PATRAKOVA, A.Ya.; TARTAKOVSKAYA, F.M.

Effect of an insulating layer on the electrical properties of
rectifiers with a TiO_2 -x base. Radiotekh. i elektron. 8
no.8:1462-1465 Aug '63. (MIRA 16:8)

1. Fiziko-tekhnicheskiy institut im. A.F.Ioffe AN SSSR.
(Electric current rectifiers)

DAVIDOVICH, N.M.; DORIN, V.A.

Electric investigation of TiO_2 -x diffusion layers on titanium.
Fiz. met. i metalloved. 16 no.2:273-277 Ag '63. (MIRA 16:8)

1. Leningradskiy fiziko-tekhnicheskii institut im. A.F. Ioffe
AN SSSR.

(Diffusion coatings—Electric properties)
(Titanium oxide)

DORIN, Y.A.; KUZNETSOV, B.I.

Device for perforating orifices in electron microscope preparation
sieves. Zav.lab. 29 no.8:1C12 '63. (MIRA 16:9)

1. Leningradskiy fiziko-tekhnicheskii institut imeni A.F. Ioffe
AN SSSR.

(Electron microscopy)

DORIN, V.A.; KOZLOV, M.M.

Measurements of potential distribution in semiconductor
rectifiers by means of a probe. Izv. vys. ucheb. zav.; fiz. no. 3:
97-101 '64. (MIRA 17:9)

1. Leningradskiy politekhnicheskij institut imeni Kalinina.

ACCESSION NR: AP4034052

S/0126/64/017/004/0536/0540

AUTHORS: Dorin, V. A.; Tartakovskaya, F. M.

TITLE: A study of the influence of oxygen generated during the reduction of TiO_2 on the oxidation of titanium

SOURCE: Fizika metallov i metallovedeniye, v. 17, no. 4, 1964, 536-540

TOPIC TAGS: titanium oxide, titanium, annealing, sodium fluoride, hydrochloric acid, oxide formation, rutile titanium

ABSTRACT: The effects of oxygen (produced by reduction of TiO_2 powder) on the physical properties of the oxide layer and on the rate of its growth were studied. It was established that it is possible to change the electrophysical properties of the TiO_{2-x} layer by immersing titanium in the oxide powder. Circular plates of Ti, 10 mm in diameter and 1.2 mm thick and with less than 0.1% impurities, were used for the oxidation experiments. Before oxidation, the plates were annealed at 1000C for 1 hour, degreased, and then pickled in an aqueous solution of 5% NaF with 12% HCl. These plates were set vertically in porcelain debitnesses and covered with TiO_2 powder pre-annealed at 800C for 3 hours. Oxidation occurred in a tubular furnace through which a constant current of steam was passed. Microphotographs of cut

Card 1/2

ACCESSION NR: AP4034052

sections were taken at 800C. Radiographic study of the structure of the material showed that the layer contained dioxide with rutile structure. As observed earlier by D. I. Layner and M. I. Tay*pin (FMM, 1960, 10, 543), the oxidation of titanium in air proceeded through molecular oxygen, whereas in steam it proceeded through atomic oxygen. The formation of atomic oxygen during the reduction of TiO_2 explained the similarities in the physical properties of the layers formed in steam and in a steam-air mixture. The increase in the contribution of atomic oxygen in the oxide layer was responsible for the growth of this layer with significant deviation in the stoichiometric properties. Orig. art. has: 5 figures, 1 formula, and 1 table.

ASSOCIATION: Leningradskiy fiziko-tokhnicheskii institut im. A. F. Ioffe AN SSSR
(Leningrad Physico-technical Institute, AN SSSR)

SUBMITTED: 28Apr63

SUB CODE: MM

NO REF SOV: 005

ENCL: 00

OTHER: 001

Card 2/2

ACCESSION NR: AP4013307

S/0032/64/030/002/0206/0206

AUTHORS: Dorin, V. A.; Kozlov, M. M.

TITLE: Silicon carbide probe for testing semiconductor materials

SOURCE: Zavodskaya laboratoriya, v. 30, no. 2, 1964, 206

TOPIC TAGS: silicon carbide, silicon carbide probe, semiconductor probe

ABSTRACT: A silicon carbide probe with a resistivity of 10 ohm-cm has been devised for testing hard semiconductor materials. The probe (see Fig. 1 of Enclosure) consists of a sharp point (1) which is fixed with tin in a copper holder (2). The holder is attached to fluorine-bearing plates (3) with a screw (4). The point scribes a line about 1.5μ wide. The contact resistance of the probe is 10^9 ohm. This instrument can be used for testing selenium and titanium dioxide. Orig. art. has: 1 figure.

ASSOCIATION: Leningradskiy politekhnicheskiy institut (Leningrad Polytechnical Institute)

Card 1/1

L 1/4976-65 EWT(m)/EWP(t)/EWP(b) Ps-4 EWT/SSD/ESD(gs) RDW/JD
 ACCESSION NR: AP5000162 8/0032/64/030/012/1190/1190

AUTHOR: Dorin, V. A.

TITLE: Charcoal-silver replicas for investigating sulfur, selenium, and tellurium

SOURCE: Zavodskaya laboratoriya, v. 30, no. 12, 1964, 1190

TOPIC TAGS: electron microscopy, surface property, sulfur, selenium, tellurium

ABSTRACT: A method is described for obtaining charcoal replicas without defects from the surfaces of sulfur, selenium, tellurium, and some of their chemical compounds. A layer of aluminum, 1 μ thick, is vaporized in a vacuum onto the surface to be investigated. Since aluminum is too brittle to be used alone, a layer of silver is applied, and further processing is the same as for the ordinary charcoal replica. Electron diffraction and electron microscope measurements show no reaction products between the layers.

ASSOCIATION: Leningradskiy politekhnicheskii institut (Leningrad Polytechnic Institute)

SUBMITTED: 00

SIB CODE: IC, NP

NO REF SOV: 000

ENCL: 00

OTHER: 000

Card 1/1

L 8555-66

ACCESSION NR: AP5021178

UR/0133/65/000/004/0112/0115

36
35
B

AUTHOR: Dorin, V. A.; Kozlov, M. M.

TITLE: The effect of defects in a p-n junction on the drift phenomenon in selenium rectifiers

SOURCE: IVUZ. Fizika, no. 4, 1965, 112-115

TOPIC TAGS: selenium rectifier, current stabilization, pn junction, rectification, electric conductivity/ TVS, AVS

ABSTRACT: The drift of selenium rectifiers TVS and AVS was investigated from the point of view of the behavior of the channels of local conductivity in a p-n junction. The fall-off of the voltage was measured for a constant current. An automatic recording potentiometer (EPP-40) with an input resistance of 10^{13} ohm was used. The current source was a battery of dry cells with a total emf of 1000 v. The voltage on the rectifier did not exceed 40 v. The investigations were carried out on pSe-nCdSe and pSe-nCdS hetero junction 40 x 40 mm in size. It was found that the resistance of a rectifier increases in time at small currents and decreases at large currents. The drifting is due to a large degree to local inhomogeneities whose conductivity varies with time. For some rectifiers there are several current values at which there is no drifting. This can be attributed to

Card 1/2

L 8555-66

ACCESSION NR: AP5021178

compensation of the loss of some conduction channels by formation of new conductivity channels. Orig. art. has: 7 figures.

ASSOCIATION: Leningradskiy politekhnicheskiy institut (Leningrad Polytechnic Institute)

SUBMITTED: 08Jan64

ENCL: 00

SUB CODE: SS, KE

NR REF SOV: 000

OTHER: 003

jw

Card 2/2

1. 27766-66 EWT(1)/EWT(m)/T/ETI/EWA(h)/EWP(t) TJP(1) JD/AT

ACC NR: AP6001590 (N) SOURCE CODE: UR/0120/65/000/006/0189/0190

AUTHOR: Dorin, V. A.

ORG: Leningrad Polytechnical Institute (Leningradskiy politekhnicheskii institut)

TITLE: A device indicating the position of the p-n junction in silicon rectifiers

SOURCE: Priory i tekhnika eksperimenta, no. 6, 1965, 189-190

TOPIC TAGS: crystal rectifier, physics laboratory instrument

ABSTRACT: A simple system is described for measuring the potential jump at the p-n junction in a thin silicon crystal plate. The transition region was measured with a precision of one micron. The p-n position was fixed by using a probe needle made of a tungsten wire. The adjustment of the needle point to the right position was made by means of the microscopes of the PMT-3 microhardness-meter and of the horizontal IZA-2 comparator. The electrical circuit included a storage battery, a recording EPP-40 electrometer with a resistance of 10^{11} to 10^{13} ohms and a d-c electrometer amplifier of Ul-2 type. The arrangement was outlined in a diagram. The distribution of potential illustrating the p-n position was shown in a curve. Orig. art. has: 2 figures.

SUB CODE: 09 / SUMM DATE: 5Nov64 / ORIG REF: 003 / OTH REF: 001

UDC: 537.33

Cord 1/1 20

L 01828-66

ENT(1)/ENT(m)/ENP(1)/ENP(m)/T/ENP(t)/ENP(b)/EMA(h) ISP(c) REM/

JD/AT

ACCESSION NR: AP5020130

UR/0109/65/010/008/1518/1522
539.293.011.41

AUTHOR: ^{49.55} Dorin, V. A.; Kozlov, M. M.

TITLE: Investigation of the potential distribution in the reverse direction in a p-Se layer adjacent to n-CdSe

SOURCE: Radiotekhnika i elektronika, v. 10, no. 8, 1965, 1518-1522

TOPIC TAGS: ^{21.17.55} selenium, cadmium selenide, electric potential, p-n junction, semiconductor device

ABSTRACT: The potential was measured at the polished end of a Se layer 80—100 μ thick. Se conductivity was 1—10 ohm-m. A thin (under 1 μ) film of n-CdSe obtained by a reactive-diffusion process formed a junction with Se. A voltage of 30—35 v was applied to the specimen, and a steel probe was set at 4 μ from the junction. Potential-distribution curves were measured. It was found that 1) the space charge at a distance of 4 μ from the junction is practically nil; 2) the reactive-diffusion-produced contact is, in fact, a defective hetero-p-n junction (conducting channels can be found in the junction); and 3) the nonuniformity of the p-n junction can be

Card 1/2

L 01828-66

ACCESSION NR: AP5020130

adequately studied by measuring the potential in Se in the reverse direction. Orig
art. has: 4 figures, 1 formula, and 1 table. [03]

ASSOCIATION: Leningradskiy politekhnicheskii institut (Leningrad Polytechnic
Institute)

SUBMITTED: 21 Jan 64

ENCL: 00

SUB CODE: 55EC

NO REF SOV: 005

OTHER: 003

ATD PRESS: 4086

Card 2/2

DAVIDOVICH, N.M.; DORIN, V.A.

Screw dislocations in diffusion layers. Fiz. met. i metalloved.
19 no.4:626-627 Ap '65. (MIRA 18:5)

1. Leningradskiy politekhnicheskoy institut imeni Kalinina.

DORIN, V.A.; KOZLOV, M.M.

Defects in p - n-junctions affecting creep in selenium rectifiers.
Izv. vys. ucheb. zav.; fiz. 8 no.4:112-115 '65. (MIRA 15:12)

1. Leningradskiy politekhnicheskiy institut. Submitted January
8, 1964.

25(1)

PHASE I BOOK EXPLOITATION SOV/1593

Dorin, Vasilii Ivanovich

Tochnaya shtampovka detaley optiko-mekhanicheskikh priborov
(Precision Cold-pressing of Parts for Optical-mechanical
Devices) Moscow, Oborongiz, 1958. 458 p. 7,050 copies
printed.

Ed.: V.Ya. Shekhter, Candidate of Technical Sciences; Ed. of
Publishing House: A.G. Kuznetsova; Tech. Ed.: V.P. Rozhin;
Managing Ed.: A.S. Zaymovskaya, Engineer.

PURPOSE: This book is intended for engineers and technicians in
optical and mechanical plants who deal with metal stampings.
It may also be used by engineers in the instrument and
watchmaking industry and by students interested in this
branch of mechanical engineering.

COVERAGE: The book deals with the cold working of metal, in
particular with the stamping of small precision parts for

Card 1/9

Precision Cold-pressing of Parts (Cont.)

SOV/1593

photographic cameras and optical equipment manufactured in the USSR. Modern methods of precision stamping on compound and progressive dies and some theoretical aspects of metal shearing are described. A new method is mentioned which is said to combine stamping and machining of parts while the parts are still in strip form. Locating parts for subsequent operations is accomplished by means of perforations along the edges of the strip. Basic forging, drawing, and upsetting operations are explained. The use of dies on an industrial scale basis, the number of dies available, and the most efficient use of these tools are discussed. Explanatory illustrations, diagrams, graphs and tables are provided. There are 52 references, of which 49 are Soviet and 3 German.

Foreword

3

Ch. I. Classification of Die Operations, Dies, and Fields of Application

5

1. Classification of cold-working operations
2. Fields of application of cold-working

5

17

Card 2/o

Precision Cold-pressing of Parts (Cont.)	SOV/1593	
Ch. II. Materials Used in Cold-Working in the Optical and Mechanical Industry		21
1. Sheet and strip steel		22
2. Aluminum and aluminum alloys		34
3. Magnesium alloys		38
4. Copper and copper alloys		39
5. Nickel and nickel alloys		45
Ch. III. Shearing Sheet Metal With Shears and Dies		48
1. Shearing with shears		48
2. Shearing with dies		50
3. Determination of forces involved in shearing on dies		52
4. Trimming operations on dies		57
5. Lay-out of parts for stamping		59
Ch. IV. Bending on Dies		65
1. Determination of the minimum bending radius		66

Card 3/9

Precision Cold-pressing of Parts (Cont.)	SOV/1593	
2. Determination of bending forces		67
3. Spring-back		68
4. Clearance between punch and die in bending		70
5. Determination of the size of the part in bending		71
6. Dies for precision bending		71
Ch. V. Drawing		77
1. Determination of the number of operations and selection of the coefficient of drawing		77
2. The use of pressure plate		80
3. Clearance (punch and die)		80
4. Punch and die corner radii		81
5. Blank for drawing		82
6. Lubrication in drawing operations		83
7. Power requirements in drawing operations		84
Ch. VI. Stamping in Compound Dies		87
1. Stamping flat parts in compound dies		89
2. Stamping shaped parts in compound dies		108
3. Trimming drawn parts in a die with horizontal or vertical movement of the punch or the die block		131

Precision Cold-pressing of Parts (Cont.)	SOV/1593	
Ch. VII. Stamping in Progressive Dies		142
1. Stamping in progressive dies of flat parts		144
2. Stamping in progressive dies of shaped parts		155
Ch. VIII. "Post-operational" Stamping With a Perforated Strip		169
1. Making flat parts		172
2. Making shaped (three dimensional) parts		178
Ch. IX. Cold Working Dies		210
1. Classification, characteristics and application of die cold working		210
2. Examples of making parts for optical and mechanical instruments with the use of die cold working		224
3. Die design for die cold working		245

Card 5/9.

Precision Cold-pressing of Parts (Cont.)	SOV/1593	
Ch. X. Some Problems of Quality Control in the Production of Precision Parts for Optical and Mechanical Instruments		
1. Organization of quality control in the shop and the instruments used		270
2. Inspection of precision parts with projectors		272
3. Preparation of scribed templates for quality control with a projector		278
4. Defects of parts produced in dies and pre- ventive measures		281
Ch. XI. Some Problems in the Design and Making of Precision Dies		286
1. Preliminary data		286
2. Shape and size of die components and determination of tolerances		289
3. Standard design of strippers and ejectors		296
4. Sectional die blocks and punches for precision dies		298
5. Making sectional die blocks and punches		304

Card 6/9

Precision Cold-pressing of Parts (Cont.)

SOV/1593

6. New methods of fastening die blocks and punches	309
7. Making die blocks and die molds by squeezing	311
8. Materials for die components	313
9. Die sets for cold working	314
Ch. XII. Organization of the Die Maintenance Services	326
1. Sequence in accepting and delivering dies	326
2. Keeping records of dies	330
3. Storing dies	332
4. Issuing to the shop, return, and checking of usability of dies	333
5. Mounting and setting-up dies on presses	335
6. Overhauling dies	339
7. Service life of dies and methods of increasing it	340
8. Safety in working with dies and presses	342

Card 7/9

Precision Cold-pressing of Parts (Cont.)	SOV/1593	
Ch. XIII. Mechanization and Automatization of Cold Working in Optical-mechanical Instrument Production		346
1. Basic trends in the development of mechanization and automation of cold working		346
2. Mechanization and automation of strip and band stock feed into the die		347
3. Mechanization of feeding separate blanks into the die		367
4. Mechanization of stripping and ejecting finished parts		382
Ch. XIV. Mechanization of Some Auxiliary Operations		391
1. Arrangement for cutting-up scrap		391
2. Machines and arrangements for cleaning operations		392
3. Machines for deburring flat parts		396
4. Mechanical sieve for separating finished parts in tumbling barrels		399

Card 8/9

Precision Cold-pressing of Parts (Cont.)

SOV/1593

Ch. XV. Equipment for Cold Working

401

1. Presses

403

2. Preparatory equipment used in cold working

445

Bibliography

455

AVAILABLE: Library of Congress

GO/rj
5-28-59

Card 9/9

DORIN, Viktor Sergeyevich; MEYLUNAS, V.P. otvetstveny redaktor;
~~MISHKOVICH, G.I.~~, redaktor; KAMOLOVA, V.M., tekhnicheskii redaktor

[How and why a ship floats] Kak i pochemu plavaet sudno.
Leningrad, Gos. soluznos izd-vo sudostroit. promyshl., 1957
113 p. (MLRA 10:5)

(Ships)

DORIN, V.S., kand. tekhn. nauk

Using differential weight equations. Sudostroenie 25 no.7:7-10
Jl '59. (MIRA 12:12)

(Naval architecture)

SEMENOV-TYANSHANSKIY, V.V., doktor tekhn.nauk; DORIN, V.S., kand.tekhn.nauk

Problems of ship reserve buoyancy and stability examined at the
Conference on the Revision of the International Convention of 1948
on the Protection of Human Life at Sea. Sudostroenie 28 no.5:1-4
My '62. (MIRA 15:7)
(Lifesaving—Congresses) (Stability of ships)

DORIN, V.S., kand. tekhn. nauk; VOLKOV, B.N., inzh.

Standardizing the reserve buoyancy of seagoing ships. Sudostroenie
28 no.5:4-12 My '62. (MIRA 15:7)
(Stability of ships)

DORIN, V.S., kand.tekhn.nauk; ARAKEL'YAN, G.V., inzh.; LOGACHEV, S.I., inzh.;
NIKOLAYEV, M.M., inzh.

Advantage of designing large-tonnage tank vessels with excess
metacentric height. Sudostroenie 29 no.7:5-8 J1 '63.
(MIRA 16:9)
(Tank vessels) (Naval architecture)

DORINOVSKAYA, A.P. [DORINOVSKAYA, A.P.]

Effect of drug-induced sleep on erythropoiesis during the post-hemorrhagic period [with summary in English] Fiziol. zhur. [Ukr.]
4 no.3:339-347 My-Je '58 (MIRA 11:7)

1. Sverdlovskiy medichniy institut, kafedra patologichnoi fiziologii.
(SLEEP)
(ERYTHROCYTES)

DORINOVSKAYA, A.P., Cand Med Sci -- (diss) "Effect of
~~energetic~~ ^{drugged} sleep ^{upon} ~~the~~ regeneration of the blood in
experimental anemia." Sverdlovsk, 1959, 15 pp (Sverdlovsk
State Med Inst) 200 copies (KL, 33-59, 121)

- 61 -

DORIS, Spalatin

Cruises with the Jadrolinija ships. Medun transp 8 no.6:416-418
Je '62.

LEVASHOVA, L.B.; DORIYENKO, Ye.P.; DEOTYAREV, V.F.

Radioactive tracer study of cobalt thiocyanate distribution between
immiscible solvents. Zhur.ob.khim.25 no.6:1066-1072 Je '55.
(MIRA 8:12)

1. Ural'skiy politekhnicheskiy institut
(Cobalt thiocyanates)

KHILINSKIY, F.A.; LOTYSHEV, I.P.; LEBEDENKO, G.B.; SHAVKUNOVA,
N.D.; DORIZO, A.P.; TERNOVAYA, K.G.; ANTIPOV, A.S.,
obshchestv. red.; BABAK, Yu.M., tekhn. red.

[Goryachiy Klyuch] Goriachii kluch. Izd.2., ispr. i
dop. [By] F.A.Khilinskii i dr. Krasnodarsk, Krasnodarskoe
knizhnoe izd-vo, 1963. 84 p. (MIRA 17:2)

1. Glavnyy vrach sanatoriya No.2 Kurorta Goryachiy Klyuch,
Kavkaz (for Lebedenko). 2. Sanatoriy No.1 Kurorta Goryachiy
Klyuch, Kavkaz (for Shavkuncva, Ternovaya). 3. Zamestitel' glavnogo
vracha po meditsinskoy chasti sanatoriya No.2 Kurorta Goryachiy
Klyuch, Kavkaz (for Dorizo).

GOYA, I., BUKUR, N., DORCA, N., RUB, D. (Bukharest)

Interrelations of rheumatic arteritis, periarteritis nodosa
and rheumatic diseases; preliminary report. Klin.med. 36
no.10:68-75 0 '58 (MIRA 11:11)

(RHEUMATISM, compl.

arteritis, relation to periarteritis nodosa (Rus))

(ARTERITIS, etiol. & pathogen.

rheum., relation to periarteritis nodosa (Rus))

(PERIARTERITIS, NODOSA

relation to rheum. arteritis (Rus))

DORKIN, Vasilii Grigor'yevich: [Dorkin, V.R.]; UKSUSOV, D. [Ukussou, D.], red.;
SLAVYANIN, I., tekhn. red.

[Selecting and training collective-farm specialists] Padbor
i vykhavanne kalhaanykh kadrau. Minsk, Dziarzh.vyd-va BSSR,
Rad.masava-palit.lit-ry, 1960. 52 p.

(Collective farms)

(Agricultural education)

(MIRA 14:3)

20904

9.6130

S/144/61/000/003/004/004
E194/E435

AUTHORS: Lifanov, V.A., Candidate of Technical Sciences, Docent,
Head of Department of Electrical Machines and Instruments
and Dorm, A.G., Senior Instructor

TITLE: An Investigation of Commutation Armature Reaction in
d.c. Machines using Hall-Effect Pick-ups

PERIODICAL: Izvestiya vysshikh uchebnykh zavedeniy,
Elektromekhanika, 1961⁴ No.3, pp.109-115

TEXT: This article describes a method of obtaining curves of
magnetic induction distribution in the air gap of a d.c. machine
using Hall-effect pick-ups. Hall-effect pick-ups used in this
work were single crystals of germanium made in the Laboratoriya
elektricheskikh mashin ChPU (Electrical Machines Laboratory of
ChPU) and their dimensions were 10 x 4 x 0.45 mm. Calibration
work showed that with inductions in the range 100 to 15000 gauss
and currents of 30 to 50 mA, the signals obtained could be
measured in an ordinary electromagnetic voltmeter without
preliminary amplification. The pick-up was used to investigate
magnetic fields in the air gap of a motor type TM-10 (PN-10);
Card 1/4

20904

S/144/61/000/003/004/004
E194/E435

An Investigation of Commutation ...

it was attached to the outer surface of one of the armature teeth. The pick-up was connected to a terminal board by flexible leads arranged so that the armature could be turned through a double pole pitch. Smooth armature rotation was ensured by a worm drive. Curves of magnetic field distribution in the air gap of d.c. machine type PN-10 are plotted in Fig.3, in which curve 1 corresponds to the armature field, curve 2 to the field of the main poles and curve 3 to the resultant field. Determination of the m.m.f. of commutational armature reaction is then considered. When the commutation in d.c. machines is accelerated or retarded there arise m.m.f.'s of commutating currents of the short-circuited armature sections which either strengthen or weaken the flux of the main poles. The fundamental theory of the effect is described in order to explain the basis of the experiments. The test circuit used is shown in Fig.4 in which the machine investigated, denoted MM in the diagram, can act as a generator or motor, it being of the same shaft as the auxiliary machine denoted BM. In this case the Hall-effect pick-up was attached under the middle of the main pole of the machine studied. The test consists in taking voltage curves at the output of the Hall-effect pick-up with the

20901

An Investigation of Commutation ... S/144/61/000/003/004/004
E194/E435

machine acting as both generator and motor. Under both conditions the speed, armature and field currents are the same. The work was done on a d.c. generator type ПН-100 (PN-100) of 115 V, 13.3 kW, 116 A, 1480 r.p.m. A single crystal germanium Hall-effect pick-up made in the Institut poluprovodnikov AN SSSR (Semiconductors Institute AS USSR) was used, its dimensions were 5 x 3 x 0.45 mm. The operating current during the test was 10 mA. The results are plotted in Fig.5. The point of intersection of the generator and motor curves corresponds to the case when the fluxes in the machines are equal in both cases. This is valid provided only that there are no m.m.f. of commutating currents, which corresponds to straight line commutation. Fig.5 also plots the difference between the generator and motor voltages and so in effect shows the change of voltage on passing from the generator to the motor conditions. This change is due to the m.m.f. of commutation currents. This m.m.f. may be determined by a special test which consists in taking a curve of the voltage on the output of the Hall-effect pick-up as function of the field current at no-load. The curve, Fig.7, is then readily constructed; it shows the relationship between the m.m.f. of commutational armature

Card 3/4

20904
S/144/61/000/003/004/004
An Investigation of Commutation ... E194/E435

reaction and the boost current in the d.c. motor. It will be seen that the boost current of 3.6 A corresponds to straight line commutation. It is concluded that this method of determining the magnetic field in the air gap of a d.c. machine is simple and easy and can be used both on the factory test bed and in teaching laboratories. The proposed method of determining the m.m.f. of commutating currents permits rapid and accurate assessment of machine commutation. There are 7 figures, 2 tables and 4 references: 3 Soviet and 1 non-Soviet.

ASSOCIATION: Kafedra elektricheskikh mashin i apparatov
Chelyabinskogo politekhnicheskogo instituta
(Department of Electrical Machines and Instruments
of the Chelyabinsk Polytechnical Institute)

SUBMITTED: October 6, 1960

Card 4/4

LIFANOV, V.A., kand.tekhn.nauk; DORM, A.G., inzh.

Use of Hall transducers for measuring the rotor angle of
synchronous machines. Vest. elektrom. 34 no.2:62-63
F '63. (MIRA 16:2)
(Electric machinery, Synchronous—Measurements)

LIFANOV, V.A., kand. tekhn. nauk, dotsent; LORM, A.G., inzh.; ROTENBERG,
M.I., inzh.

Method for the automatic synchronization of synchronous
machines. Izv. vys. ucheb. zav.; energ. 7 no.10:84-87
0 '64. (MIRA 17:12)

1. Chelyabinskiy politekhnicheskii institut.

L 01132-66

ACCESSION NR: AP5017467

UR/0144/65/000/006/0718/0720
621.313.33+621.3.047

AUTHOR: Lifanov, V. A. (Docent); Dorm, A. G. (Senior lecturer)

27
B

TITLE: Measuring and oscillographing the slip in induction machines

SOURCE: IVUZ. Elektromekhanika, no. 6, 1965, 718-720

TOPIC TAGS: induction machine, slip

ABSTRACT: A method of measuring the slip of an induction machine by means of a commutator-type tachometer generator mechanically coupled to the machine is suggested. The tachogenerator stator has distributed 3-phase winding whose number of poles equals to that of the main machine; both are connected to the same a-c supply. EMF across the tachogenerator brushes, directly proportional to the slip of the induction machine, can be easily measured by an oscillograph. Oscillograms of the slip of a 3.6-kw, 380-v, 2890-rpm induction motor when the rated load was suddenly thrown on (or varied) are presented. An auxiliary use of a Hall generator is also suggested. Orig. art. has: 6 figures and 6 formulas.

Card 1/2

L 01132-66

ACCESSION NR: AP5017467

ASSOCIATION: Chelyabinskiy politekhnicheskiy institut (Chelyabinsk Polytechnic Institute)

SUBMITTED: 28Nov62

ENCL: 00

SUB CODE: EE

NO REF SOV: 001

OTHER: 00

Card 2/2

DP

BIRULYA, I.N., kand.tekhn.nauk, dotsent; DORM, F.A., inzh.

Adjustment and study of the commutation of large d.c. generators.
Energ. sbor. no.2:3-12 '59. (MIRA 15:1)

(Electric generators)
(Commutation (Electricity))

DORMAKOVICH, Petr Andreyevich; MIKHALKOV, Aleksandr Vladimirovich;
PETROV, Aleksandr Vasil'yevich; POYARKOV, K.M., red.;
BORUNOV, N.I., tekhn. red.

[Manufacture and maintenance of gas-discharge light fixtures]
Izgotovlenie i obsluzhivanie gazosvetnykh ustanovok. Moskva,
Gosenergoizdat, 1962. 54 p. (Biblioteka elektromontera, no.72)
(MIRA 15:12)

(Fluorescent lamps) (Fluorescent lighting)

DORMAKOVICH, P.A.

Methods for detecting faults in gas-discharge lamps in commercial
signs. Prom. energ. 18 no.11:60 N '63. (MIRA 16:12)

SHPARBER, L.Ya.; DORMAN, A.I.

Ladle travel in one-lip casting of iron. Metallurg 5
no. 12:10-12 D '60. (MIRA 13:11)

1. Magnitogorskiy metallurgicheskiy kombinat.
(Blast furnaces—Equipment and supplies)

DORMAN, A.I.; LESHCHINSKIY, L.Z.; KIYASHKO, V.S.; BAKSHINOV, A.S.;
LUKASHOVA, A.N.

Pneumatic delivery of specimens of cast iron, steel, and slag
to the chemical laboratory. Metallurg 9 no.10:12-13 0 '64
(MIRA 18:1)

1. Magnitogorskiy metallurgicheskiy kombinat.

ROGANOV, Boris Ivanovich, doktor tekhn. nauk [deceased]; DZHABAROV, Gafar Dzhabarovich, kand. tekhn. nauk; KOTOV, Dmitriy Andreyevich, kand. tekhn. nauk; BALTABAYEV, Sultan Dusayevich, kand. tekhn. nauk; SOLOV'YEV, Nikolay Dmitriyevich, inzh.; DORMAN, I.M., retsenzent; DUKHOVNIY, F.N., red.; SOKOLOVA, V.Ye., red.

[Primary processing of cotton] Pervichnaia obrabotka khlopka.
[By] B.I.Roganov i dr. Moskva, Legkaia industriia, 1965.
485 p. (MIRA 18:12)

L 65298-65 EWT(l)/EWT(m)/FCC/T/EWA(h) IJP(c) CW

ACCESSION NR: AP5020995

UN/0203/65/005/004/0666/0672

523.165

57

54

B

AUTHORS: Dorman, I. V.; Dorman, L. I.

TITLE: Proton and α -particle modulation in regions of small intensity and the cosmic ray spectrum in the galaxy /9

SOURCE: Geomagnetizma i aeronomiya, v. 5, no. 4, 1965, 666-672

TOPIC TAGS: proton, alpha particle, cosmic ray, galaxy, beam modulation, spectrum analysis 12,44,55 12,55

ABSTRACT: From alpha particle and proton spectra the governing parameters of the modulation mechanism are investigated, which in turn sheds some light on the intensity gradients of cosmic rays from interplanetary space. The study is carried out in three parts. In part I, the 11-year variation in cosmic rays and the modulation of proton and α -particle beams are analyzed in the low energy range. The modulated primary cosmic ray spectra are represented by

which, when combined with the velocity ratio of protons to α -particles, yields

$$\frac{D_{\text{mod}, p}(R)}{D_{\text{mod}, \alpha}(R)} = K \exp \left[\frac{a(R)}{v_p} \left(\sqrt{\frac{R^2 + 4}{R^2 + 1}} - 1 \right) \right],$$

Card 1/2

L 65298-65

ACCESSION NR: AP5020995

where R is the ray intensity (hardness). From this, the transport path for scattering of 1 Bev particles at $r_{max} \sim 100$ is calculated to be 2.10^{11} cm. In part II, the spectra of galactic cosmic rays at low energy are analyzed during minimum solar activity. It is shown that the famous "link" observed in the cosmic ray spectra is wholly due to modulation of cosmic rays in interplanetary space. Finally, in part III, the above results are used to calculate the expected cosmic ray intensity gradients during July, 1963 for particle intensities of 1 and 0.7 Bev. These yield 3.3%/1 a.e. for protons and 5.3%/1 a.e. for α -particles in the first case, and 4.1%/1 a.e. for protons and 7%/1 a.e. for α -particles in the second case. Orig. art. has: 13 formulas, 1 table, and 1 figure.

ASSOCIATION: Institut zemnogo magnetizma, ionosfery i rasprostraneniya radiovoln, AN SSSR (Institute of Terrestrial Magnetism, Ionosphere, and Radio Wave Propagation, AN SSSR)

44,55

SUBMITTED: 05Oct64

ENCL: 00

SUB CODE: 02

NO REF SOV: 006

OTHER: 009

Card 2/2

L 16028-66 ENT(1)/EWA(h)/FCC GW

ACC NR: AT6003522

SOURCE CODE: UR/3184/65/000/007/0005/0017

AUTHOR: Dorman, I. V.; Dorman, L. I. (Dr. of Physico-Mathematical Sciences)

30
B+1

ORG: none

TITLE: Investigation of the 11-year cosmic ray cycle (from data of observations at sea level)

SOURCE: AN SSSR. Mezhdunarodstvennyy geofizicheskiy komitet. Kosmicheskiye luchy,
no. 7, 1965, 5-17

TOPIC TAGS: cosmic ray, geomagnetism, sunspot cycle, magnetic rigidity

ABSTRACT: Data from a world-wide network of neutron monitors, measurements of latitude effects in the neutron and hard components of cosmic rays at sea level, and stratospheric measurements of proton and α -particle fluxes in the low energy region made from high altitude balloons are used as a basis for analyzing the 11-year cosmic ray cycle. It is found that the amplitude of the 11-year variation increases with a reduction in the geomagnetic cutoff hardness R which indicates that the primary spectrum of the 11-year variation decreases with an increase in R . The lag in

Card 1/2

2

2

L 16028-66

ACC NR: AT6003522

changes of cosmic ray intensity with respect to changes in solar activity increases with a reduction in the penetrating power of the particles. There is a close relationship between cosmic ray intensity, the number of sunspots and radiation flux at 10.7 cm. When considering variations for shorter periods of time, there is a closer relationship between the intensity of cosmic rays and the K_p -index of magnetic activity, while longer time intervals show a closer correlation between intensity and the number of sunspots. Hysteresis phenomena are studied on the basis of neutron monitor data and observations in the stratosphere. For $R > 3$ Bev the energy spectrum for the 11-year cycle is given in the form $\delta D(R)/D(R) \sim R^{-(0.8-0.9)}$ which is steeper for $R > 13$ Bev with $\gamma > 2$. The spectrum of the 11-year cycle was considered together with delay phenomena to determine the scattering path as a function of particle hardness and the dimensions of the modulating space. The magnetic field intensity in scattering nonhomogeneities is determined for various assumptions on the relationship between the average dimensions of nonhomogeneities and the distance between them. The contribution of variability in the lag effect is evaluated. Orig. art. has: 8 figures, 16 formulas.

SUB CODE: 08/ SUBM DATE: 00/ ORIG REF: 013/ OTH REF: 019

Card 2/2

USSR.

(1953). The influence of meteorological factors on the latitude effect of the hard component of cosmic rays and the process of the generation of mesons.

DORMAN, Letter in Zh. Eksp. teory. Fiz., 26, No. 4, 1953-1954 in Russian.

The author's recent work (Dokl. Akad. Nauk, SSSR, 94, No. 3, 433; 95, No. 1, 49 (1954)) makes it possible to correct the latitude effect for the increase of intensity associated with the temperature variation R_m in the equator to $\lambda = 55^\circ$ (about 4%). The corrected intensity levels off very clearly between 35° - 40° , corresponding to a minimum primary proton energy of 6.8×10^9 eV. This is interpreted as meaning that protons with less energy cannot generate μ -mesons with energy $> 2 \times 10^9$ eV, necessary for their detection at sea level.

W. J. SWIATECKI

Sci Res Inst. Terrestrial Magnetism

DERMAN, L.I.
USSR/Nuclear Physics - Cosmic rays in meteorology

Card 1/1

Pub. 147-3/16

Author

: Dorman, L. I.; Kuz'min, A. I.; Tyanutova, G. V.; Feynberg, Ye. L.;
Shafer, Ya. G.

Title

: Variations in the intensity of cosmic rays and the role of meteorological factor

Periodical

: Zhur. eksp. i teor. fiz. 26, 537-544, May 1954

Abstract

: Briefly expound the results of an experimental and theoretical study of the influence of meteorological factors on the observed (at sea level) intensity of the hard component of cosmic rays. Show that knowing the distribution of temperature in the atmosphere above the observation point one can allow for the meteorological factors with an accuracy up to 0.1 to 0.2% in the intensity of cosmic rays. Here the remaining divergence lies within the limits of error of the given meteorological sounding. It turns out that the seasonal variations in the intensity of the hard component are due to meteorological factors. The daily variations are essentially masked by these factors.

Submitted

: October 27, 1953

"On the Theory of Meteorological Effects on Cosmic Radiation."
Dokl. Akad. Nauk SSSR, V. 94, No. 3, 433-6, 1954.

An analytical expression relating the μ -meson intensity at a given height to various meteorological factors. is deduced.

Sci. Res. Inst. Terrestrial Magnetism.

LOHMAN, L. I.

"On the Temperature Effect of the Hard Component of Cosmic Rays."
Dokl. Akad. Nauk SSSR, Vol. 95, No. 1, 49-52, 1954.

Duperier's theory of this effect does not, in general, lead to values of
the temperature coefficient which are in agreement with experiment.

DORMAN, L. I.; KAMINER, N. S.; KOYAVA, V. K., SHAFER, Yu. G. and SHVARTSMAN, B. F.

"Observation of the Large Cosmic Ray Increase of February 23, 1956
in the USSR."

✓ Scientific Research Institute of Terrestrial Magnetism (Moscow)
Sverdlovsk Geophysical Observatory
Tbilisi Cosmic Ray Station
Yakutsk Affiliate of the Academy of Sciences of the USSR
Cape Schmidt Cosmic Ray Station

Nuclear Physics, 1. No. 8, 1956, p. 585-592

9402

ON THE ORIGIN OF VARIATIONS IN COSMIC RAYS. I

Durham (Research Inst. of Earth Magnetism)

1950-51

The method of coupling coefficient ϵ and a new

evaluation theory of meteorological effects

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

of the magnetic field

HP Ditz

DORMAN, L. I.

"Unusual Surge of Cosmic Ray Intensities." Priroda, 45, 85-7, 1956


An unusual surge of cosmic-ray intensity was observed Feb. 23 1956. The scale showing the distribution of the cosmic-ray intensities indicate stronger concentration of intensities at higher latitudes (Moscow) and considerable lowering of intensities at the lower latitudes (Japan). The observed surge was 5 to 10 times stronger than the ones observed in 1942 or 1946. The surge of cosmic rays was combined with powerful chromospheric sun eruption and interrupted all short-wave radio communications. It is assumed that the eruption took place on the northwest tip of the sun where a great deal of activity, large number of spots, and chromospheric eruptions were observed before and after the event.

Research Inst. of Earth Magnetism, Ionosphere and Distribution of Radiowaves.

DORMAN, L. I. and FEYNBERG, Ye. L.

"Variations in Intensities of Cosmic Rays." Uspekhi Fiz. Nauk 59, 189-228, 1956.

Variations of cosmic-ray intensities caused by the atmospheric conditions and by acceleration, slowing down, and scattering in the corpuscular streams emitted by the sun, and by large and small sun eruptions are reviewed. Analyses of various phenomena and methods of study are discussed.



DORMAN, Leyb Isaakovich; GRIGOROVA, V.A., redaktor; AKHLAMOV, S.N., tekhnicheskii redaktor

[Variations in cosmic rays] Variatsii kosmicheskikh luchei. Moskva, Gos.izd-vo tekhniko-teoret.lit-ry, 1957. 492 p. (MIRA 10:8)
(Cosmic rays)

83800

S/035/59/000/003/006/039/
A001/A001

3.1800

Translation from: Referativnyy zhurnal, Astronomiya i Geodeziya, 1959, No. 3,
pp. 30-31, # 1902

AUTHOR: Dorman, L. I.

TITLE: Information on Solar Corpuscular Fluxes, Obtained from Studying
Variations in Cosmic Radiation 19

PERIODICAL: V sb.: Fiz. solnechn. korpuskulyarn. potokov i ikh vozdeystviye
na verkhnyuyu atmosferu Zemli, Moscow, AN SSSR, 1957, pp. 112-125,
Discuss. 125-128

TEXT: The author points out that the study of variations in cosmic
radiation makes it possible to obtain information on physical conditions in the
✓ interstellar and interplanetary media, on processes on the Sun, and other data
of astrophysical nature. The connection between astrophysical and geophysical
phenomena (flares on the Sun, magnetic storms, 11-year solar activity cycles,
etc.) on the one hand, and changes in the intensity of cosmic radiation, on the
other hand, indicates a great role of the Sun in the origin of cosmic rays. X

and 1/2

83800

S/035/59/000/003/006/039

A001/A001

Information on Solar Corpuscular Fluxes, Obtained from Studying Variations in Cosmic Radiation

However, distortions introduced in the observed variations by changes of meteorological conditions and by contributions of secondary particles originated in the atmosphere, make it difficult to discover the true connection between variations of cosmic radiation and astrophysical factors. At present a method has been developed of excluding meteorological factors. Moreover, a method of "connection coefficients" has been developed which makes it possible to determine variations of the primary flux from the observed variations in secondary components of cosmic rays. The performed analysis of solar-diurnal variations of cosmic radiation, connected with the geomagnetic and solar activities, and its connection with corpuscular fluxes warrant some conclusions on the nature of these fluxes. There exist two types of corpuscular fluxes. The first type fluxes carry comparatively weak magnetic fields ($\sim 10^{-5}$ erg near the Earth), are of low density (a few particles per 1 cm^3 near the Earth) and induce weak and moderate magnetic disturbances on hitting the Earth; on an average, 5-10 fluxes are ejected simultaneously, and this number depends only slightly on the solar activity. The fluxes of this type are due to high-latitude formations on

Cont. 2/2

83800

S/035/59/000/003/006/039
A001/A001

Information on Solar Corpuscular Fluxes, Obtained from Studying Variations in
Cosmic Radiation

the Sun and are ejected approximately symmetrically with respect to the solar equator. The solar-diurnal variations are connected mainly with the cosmic rays of this type. In the fluxes of the second type the intensity of magnetic field is higher than 10^{-4} - 2×10^{-4} erg, density $\sim 10^3$ cm $^{-3}$; they give rise to intensive and very intensive magnetic storms on the Earth. The number of second-type fluxes ejected by the Sun is well correlated with Wolf numbers and varies considerably during the cycle of solar activity; these fluxes are connected with low-latitude formations (possibly with sunspots). 27-day variations of cosmic rays, reductions of intensity during magnetic storms, dependence on solar activity, etc. are due to the fluxes of the second type. There are 30 references.

M. I. Fradkin

Translator's note: This is the full translation of the original Russian abstract.

Card 3/3

DORMAN 17

AUTHOR: Blokh, Ya. L. and Dorman, L. I.

120-2-13/37

TITLE: Meteorological Coefficients for 4π and 2π Counter Telescopes. (Meteorologicheskiye Koeffitsiyenty dlya Kubicheskogo i Polukubicheskogo Teleskopov.)

PERIODICAL: Priory i Tekhnika Eksperimenta, 1957, No.2, pp. 46 - 48 (USSR).

ABSTRACT: Complex telescopes for continuous registration of the cosmic rays intensity will be extensively used during the coming I.G.Y. The author determines the theoretical values of the barometric pressure coefficient and of the temperature coefficient which can be used with cubical telescopes at the sea level and to the semi-cubical telescopes under the surface of the earth, at depths of 25 and 55m of the water equivalent. The use of these coefficients will free experimental data from the distorting influence of the varying meteorological conditions. According to the theory (Ref. 1) the relative change

$$\frac{\delta N_{\mu}}{N_{\mu}} = \alpha_{\text{bar}} \delta h_0 + \int_0^h W_t(h) \cdot \delta T(h) dh,$$

of μ -mesons intensity due to a relative barometric change Card 1/3 δh_0 at the observation level and the relative temperature

120-2-13/37

Meteorological coefficients for 4π and 2π Counter Telescopes.

change $\delta T(h)$ is given by eq.(1). This expression has two coefficients a_{bar} - atmospheric pressure coefficient in %/millibar and $w_T(h)$ - density of the temperature coefficient in %/1°C atm. Both coefficients can be determined from the two equations (2), where $w_T(h, \epsilon)$ and $a_{bar}(\epsilon)$ are the meteorological coefficients for directed intensities. The values of $w_T(h, \epsilon)$ for subterranean measurements were obtained in Ref.2 and for the sea level measurements in Ref.3. Results of a calculation of $a_{bar}(\epsilon)$ are given in the form of a graph (Fig.2) for various assumed values of the effective index n of the differential spectrum of π -meson production. Using $a_{bar}(\epsilon)$ and $w_T(h, \epsilon)$ as defined in eq.(2), the required coefficients are presented graphically in Figs. 3 and 4 (cf. Ref.2). The meteorological coefficients should not depend in practice on the latitude of the observation point, nor on the local climatic conditions. Six graphs of numerical

Card 2/3

120-2-13/37
Meteorological coefficients for 4π and 2π Counter Telescopes.
reference are given. There are 7 references, 5 of which
are Slavic.

SUBMITTED: November 3, 1956.

ASSOCIATION: Scientific and Research Institute of Terrestrial
Magnetism, of the Ionosphere and of Radiowave Propagations.
(Nauchno-Issledovatel'skiy Institut Zemnogo Magnetizma,
Ionosfery i Rasprostraneniya Radiovoln.)

AVAILABLE: Library of Congress.

Card 3/3

DORMAN, L.I.

see card 4-6

PHASE I BOOK EXPLOITATION 881

Akademiya nauk SSSR. Yakutskiy filial

Variatsii intensivnosti kosmicheskikh luchey (Variations of the Intensity of Cosmic Rays) Moscow, Izd-vo AN SSSR, 1958. 168 p. (Series: Its: Trudy, seriya fizicheskaya, vyp. 2) 1,500 copies printed.

Resp. Ed.: Shafer, Yu.G., Candidate of Physical and Mathematical Sciences; Ed. of Publishing House: Fradkin, M.I.; Tech. Ed.: Pavlovskiy, A.

PURPOSE: This collection of articles is for scientists and students of cosmic rays and meteorology.

COVERAGE: This issue contains articles on experimental methods in the continuous registration of cosmic rays, the investigation of meteorological effects of the different components of cosmic rays, and the connection between variations in cosmic ray intensity and solar and magnetic activity. Part I describes apparatus used in

Card 1/6

Variations of the Intensity of Cosmic Rays 881

measuring cosmic ray intensity on and under the earth's surface and in the upper layers of the atmosphere, and specifically discusses the ASK automatic ionization chamber. Part II discusses the theory, methods and results of the investigation of meteorological effects of the various components of cosmic rays. Part III discusses the characteristics of daily variations in cosmic ray activity. The following scientists are mentioned in the introduction: S.N.Vernov, Corresponding Member of the AS USSR, Professor Ye.L.Feynberg, and N.L.Grigorov, Doctor of Physical and Mathematical Sciences. The articles are accompanied by diagrams, tables, and bibliographic references.

TABLE OF CONTENTS:

Preface

3

Card 2/6

. Variations of the Intensity of Cosmic Rays 881

PART I. APPARATUS FOR MEASURING VARIATIONS
OF INTENSITY OF COSMIC RAYS

Shafer, Yu.G. Continuous Registration of Variations in the Intensity
of Cosmic Rays by an Ionization Chamber With Automatic Control 7

Shafer, Yu.G. Further Improvements in Automatic Cosmic Ray Stations 23

Kuz'min, A.I., Scripin, G.B., Varygin, A.V., Installation for
Studying the Energy Characteristics of Cosmic Ray Variations 34

Kuz'min, A.I., Varygin, A.V. Apparatus for Subsurface Measurement
of Variations in Cosmic Ray Intensity 36

Belomestnykh, V.A., Shafer, Yu.G. Methods of Registration and
Study of Cosmic Ray Intensity Variations in the Stratosphere 47

Card 3/6

- Variations of the Intensity of Cosmic Rays 881

PART II. METEOROLOGICAL EFFECTS

- Dorman, L.I. Concerning the Theory of Meteorological Effects of the General Ionizing and the Soft Components of Cosmic Rays 59
- Dorman, L.I. Concerning the Theory of Temperature Effect of the Neutron Component in Cosmic Rays 68
- ✓ Kaminer, N.S. Problem of Temperature Extrapolation in the Lower Stratosphere Region 73
- ✓ Sokolov, V.D. Methodology of Applying Temperature Corrections in the Intensity Measurements of the Hard Component Cosmic Rays Using the Atmospheric Temperature Profile up to Elevations of 5-6 km 78
- ✓ Tyanutova, G.V. Preliminary Results of Comparing Data on Measurement of Global Intensity Variation of the Hard Component of Cosmic Rays with Instruments ASK-1 and S-2 81

Card 4/6

Variations of the Intensity of Cosmic Rays 881

Koval'skaya, A.I. Problem of Seasonal Variability of the Barometric Coefficient of the Hard Component of Cosmic Rays 85

Koval'skaya, A.I., Krasil'nikov, D.D., Nikol'skiy, S.I. Preliminary Results in Determining the Barometric and Temperature Effects of Extensive Torrential Rains Near the Sea Level 88

PART III. NATURE OF VARIATIONS IN COSMIC RAY INTENSITY

Glokova, E.S., Kaminer, N.S., Mishina, N.A. Cyclic and Seasonal Variations of Diurnal Wave of Intensity of Cosmic Rays 95

Kuz'min, A.I., Skripin, G.V. Relation of the Diurnal Effect of Cosmic Ray Intensity to Geomagnetic and Solar Activity 107

Fuks, L.A., Shvartsman, B.F. Temperature Effect in Seasonal and Diurnal Variations of the Hard Component of Cosmic Rays From the Data Collected on Shmidt's Promontory Station 118

Card 5/6

Variations of the Intensity of Cosmic Rays 881

Sokolov, V.D. On the Nature of 27-day Variations in Cosmic Ray Intensity 123

~~Dorman, L.I., Freidman, G.I. Interpretation of the Cosmic Ray Burst of February 23, 1956~~ 129

List of Symbols Used 170

AVAILABLE: Library of Congress

MM/whl
1-21-59

Card 6/6

DORMAN, L.I.

P.3

PHASE I BOOK EXPLOITATION

80V/3762

Konferentsiya po magnitnoy gidrodinamike. Riga, 1958.

Voprosy magnitnoy gidrodinamiki i dinamiki plazmy; trudy Konferentsii.
(Problems in Magnetohydrodynamics and Plasma Dynamics; Transactions of a
Conference) Riga, Izd-vo AN Latvyskoy SSR, 1959. 343 p.
Errata slip inserted. 1,000 copies printed.

Sponsoring Agency: Akademiya nauk Latvyskoy SSR. Institut fiziki.

Editorial Board: D.A. Frank-Kamenetskiy, Doctor of Physics and Mathematics,
Professor; A.I. Vol'dek, Doctor of Technical Sciences, Professor; I.M. Kirko,
Doctor of Physics and Mathematics; V.Ya. Veldre, Candidate of Physics and
Mathematics; V.G. Vitol, Candidate of Physics and Mathematics; Yu.M. Krumin';
and V.Ya. Kravchenko.

Ed.: A. Teytel'baum; Tech. Ed.: A. Klyavinya

PURPOSE: This book is intended for physicists working in the field of magneto-
hydrodynamics and plasma dynamics.

Card 1/12

5

Problems in Magnetohydrodynamics(Cont.)

NOV/3762

COVERAGE: This volume contains the transactions of a conference held in Riga, June 1958, on problems in applied and theoretical magnetohydrodynamics. The objects of the conference were the investigation of the basic trends in theoretical and applied magnetohydrodynamics, establishing contact between the people doing research in different branches of magnetohydrodynamics, and promoting the participation of theoretical physicists in problems in applied magnetohydrodynamics. More than 160 persons from different parts of the Soviet Union took part in the conference, and 55 papers were read. Similar conferences are to be held regularly in the future; the next such conference is scheduled to be held in Riga in June 1960. In this present collection of the transactions of the conference, most of the papers and comments on papers are presented by the authors themselves in an abridged form. The book is divided into two parts: the first part deals with problems in theoretical magnetohydrodynamics and plasma dynamics, and consists of 35 articles on such aspects of the problem as the application of magnetohydrodynamics in astrophysics (D.A. Frank-Kamenetskiy), magnetohydrodynamics and the investigation of cosmic-ray variations (L.I. Dorman), acceleration of plasma in a magnetic field (G.V. Gordeyev and A.I. Gubanov), stability of shock waves and magnetohydrodynamics (A.I. Akhiezer.). The second part, consisting of 33 articles, deals with problems of experimental magnetohydrodynamics, including the application of physical simulation for investigation of electromagnetic processes in liquid metals (I.M. Kirko) and the development of electromagnetic pumps (P.G. Kirillov), at the Institute of Physics of the

Card 2/10

Problems in Magnetohydrodynamics (Cont)

80V/3762

Academy of Sciences, Latvian SSR. Several articles are devoted to induction pumps, electromagnetic crucibles, electromagnetic stirrers for molten metals, and their application in the metallurgical industry including schematic diagrams of their power-supply systems. References are given at the end of most of the articles.

TABLE OF CONTENTS:

Foreword	3
PROBLEMS IN THEORETICAL MAGNETOHYDRODYNAMICS AND PLASMA DYNAMICS	
Frank-Kamenetskiy, D.A. The Role of Magnetohydrodynamics and Plasma Dynamics in Certain Problems in Astrophysics	7
<u>Dorman, L.I.</u> Magnetohydrodynamics and Research in Cosmic-Ray Variations	13
Syrovatskiy, S.I. The Cosmic-Ray Spectrum and the Significance of Cosmic Rays in Cosmic Gasdynamics	45
Card 3/10 5	

Problems in Magnetohydrodynamics (Cont)

SOV/3762

Velikhov, Ye.P. The Influence of a Magnetic Field on the Flow Stability of a Conducting Fluid	49
Terletskiy, Ya.P. Certain Problems of the Movement of Rarefied Plasma in a Magnetic Field	59
Sagdeyev, R.Z. On Nonlinear Steady Flow of Rarefied Plasma in a Magnetic Field	63
Braginskiy, S.I. One Criterion for the Applicability of Magneto-hydrodynamic Equations to Plasma	67
Polovin, R.V. Comments on the Paper	71
Gordeyev, G.V., and A.I. Gubanov. The Problem of Plasma Acceleration in a Magnetic Field	73
Gordeyev, G.V. Comments on the Paper	74
Dorman, L.I., and G.I. Freydmann. On the Possibility of Charged-Particle Acceleration by Shock Waves in Magnetized Plasma	77

Card 4/22

Problems in Magnetohydrodynamics(C ont)

80V/3762

Dorman, L.I. On Charged-Particle Acceleration During Powerful
Impulse Discharges and the Collision of Magnetized Clouds

83

Konyukov, M.V. The Effect of Longitudinal Magnetic Fields on
Electron Temperature in Plasma

89

Kholev, S.R. Research on Certain Characteristics of Plasma of
Xenon and Argon Behind a Powerful Shock Wave

93

Granovskiy, V.L., K.P. Ryumina, V.I. Savoskin, and G.G. Timofeyeva.
Investigation of an Electrodynamically Pinched Arc With the Aid of
an Electron-Optical Converter

107

Akhlyezser, A.I., G.Ya. Lyubarskiy, and R.V. Polovin. On the Stabili-
ty of Shock Waves in Magnetohydrodynamics

116

Kontorovich, V.M. On the Interaction of Small Disturbances With
Discontinuities and the Stability of Shock Waves in Magnetohydrodynamics

117

Card 5/12
5

Doc man, 61

- Report on the 31st Conference on Magnetohydrodynamics, 1979, 2-10 July 1979, Moscow, 1979, 317 pp.
- 1979, 2-10 July 1979, Moscow, 1979, 317 pp.
- The subject of the 31st conference reports and discussions of reports are presented in the source in abbreviated form. Previously published reports are included there as brief abstracts only. The material published there for the first time (abbreviated and unabbreviated) are as follows:
- "The Role of Magnetohydrodynamics and Plasma Dynamics in Cosmic Problems of Astrophysics," by B. A. Piontchik, Moscow, pp 7-13.
- "Magnetohydrodynamics and the Study of Variations of Cosmic Rays," by L. I. Dorman, Moscow, pp 13-14.
- "Cosmic Ray Spectra and Their Role in Cosmic Ray Dynamics," by S. I. Ginzburg, Moscow, pp 15-16.
- "The Influence of a Magnetic Field on the Stability of Flow of a Conducting Fluid," by V. A. Il'inskiy, Moscow, pp 16-18.
- "Some Problems of the Motion of a Barred Plasma in a Magnetic Field," by V. A. Il'inskiy, Moscow, pp 19-20.
- "On Solitary Steady-State Motion of a Barred Plasma in a Magnetic Field," by V. A. Il'inskiy, Moscow, pp 21-22.
- "On the Criterion of Applicability of the Equations of Magnetohydrodynamics to a Plasma," by S. I. Ginzburg, Moscow, pp 23-24. (Discussion of the report by V. A. Il'inskiy, Moscow, pp 25-26).
- "On the Possibility of Accelerating Charged Particles by Means of Shock Waves in a Magnetized Plasma," by V. A. Il'inskiy and G. I. Il'inskiy, Moscow, pp 27-28.
- "On an Acceleration of Charged Particles During Powerful Impulse Discharges and During the Collision of Magnetic Clouds," by L. I. Dorman, Moscow, pp 29-30.
- "The Influence of a Longitudinal Magnetic Field on the Temperature of the Electrons in a Plasma," by N. V. Kopylov, Tula, pp 31-32.
- "Temperatures of Cosmic Characteristics of a Plasma of Iron and Argon Behind a Powerful Shock Wave," by N. V. Kopylov, Moscow, pp 33-34.
- "Observation of Electrodynamical Contractions of an Arc with the Aid of an Electron-Optical Converter," by L. L. Ginzburg, K. A. Ginzburg, Z. A. Ginzburg, and G. G. Ginzburg, Moscow, pp 35-36.
- "On the Interaction of Weak Perturbations with Discontinuities and the Stability of Shock Waves in Magnetohydrodynamics," by V. M. Kuvshinov, Moscow, pp 37-38.
- "On the Stability of Shock Waves in Magnetohydrodynamics," by V. M. Kuvshinov, Moscow, pp 39-40.
- "On the Scattering of Hydromagnetic Waves on Turbulent Fluctuations," by A. G. Sivukhin and Yu. A. Kuvshinov, Moscow, pp 41-42.
- "On the Damping of Magnetohydrodynamic Waves in a Plasma," by S. I. Ginzburg, Moscow, pp 43-44.
- "Single Wave in Magnetohydrodynamics," by A. I. Akhiezer, G. Ya. Akhiezer, and R. V. Akhiezer, Moscow, pp 45-46.
- "Two-Dimensional Problems of Magnetohydrodynamics," by G. B. Ginzburg, Moscow, pp 47-48.
- "On Wave-Induced Flows in Magnetohydrodynamics," by A. I. Akhiezer, Moscow, pp 49-50.
- "Oscillations of an Infinite Gas Cylinder with its Own Gravitation in a Magnetic Field," by L. N. Tsimmerman, Moscow, pp 51-52.
- "On Magnetic Boundary Layers and Electric Current Discharges in Moving Media," by V. A. Il'inskiy, Moscow, pp 53-54.

DORMAN L. I.

"CONCERNING THE ENERGY SPECTRUM OF THE VARIATIONS AND DURATION OF THE INCREASE EFFECT PRECEDING MAGNETIC STORMS"

L. I. Dorman

To explain the effect of cosmic ray intensity increases preceding magnetic storms, a mechanism of accelerating particles by a shock wave formed by the front edge of the solar corpuscular stream is suggested. The shock wave is propagated in interplanetary space with a velocity of approximately 1.3 times that of the stream and reaches the Earth earlier than the stream. It is therefore a sort of "forerunner" of the magnetic storm.

Calculations are made of the energy spectrum of the particles evoking the increase effect and of the duration of this effect. It is shown that with decrease in rigidity of particles the amplitude of this effect should increase and the duration should decrease. Thus, the expected duration of the effect for particles with a rigidity of $3 \cdot 10^{10}$ ev/c is approximately 8 hours; it is about 20 minutes for particles with a rigidity of 10^9 ev/c, and approximately 2 minutes for particles with a rigidity of 10^8 ev/c (in the latter case, the amplitude of the effect should be quite appreciable and approach 100%). By means of correlation coefficients, calculations are made of the expected variations of different components at different latitudes. A preliminary comparison of the obtained results and the experimental data is made, and other possible explanations of this effect are discussed.

report presented at the International Cosmic Ray Conference, Moscow, 6-11 July 1959

DORMAN L.I.

"THE POSSIBILITY OF PRE-ACCELERATION OF COSMIC RAYS BY THE PINCH EFFECT
MECHANISM"
L.I. Dorman

A model of charged particle acceleration between two approaching half-spaces with "frozen" magnetic fields is considered. Assuming the absence of magnetic field between the half spaces, the law of energy change with time is determined, as well as the maximum relative acceleration as a function of initial particle momentum, the distance between the half spaces and the intensity of the "frozen" magnetic field.

An evaluation is made of the conditions of injection, the acceleration of particles with different Z is compared and the energy spectrum for the acceleration from thermal energies with initial Maxwell distribution is determined. The variations in the results are evaluated for the case when the discussed model is generalized to cover real magnetic cloud collisions and pinch effect in the plasma. On the basis of A. B. Severny's observations, and on results, the significance of the mechanism considered is discussed for pre-acceleration of particles in solar flares and supernovae and also for collisions of magnetized clouds (in particular for collisions of galaxies). The assumption is made that further acceleration of particles to higher energies is accomplished by Fermi's statistical mechanism.

report presented at the International Cosmic Ray Conference, Moscow 6-11 July 1959